

		7-10 Years Old	11 Years and Older
HepB Hepatitis B	3 doses The final dose of HepB must be given at 24 weeks of age or older. Only 3 doses are required if the 3 <sup>rd</sup> dose was received at or after the child was 24 weeks of age; otherwise 4 doses are required.		
Polio Poliomyelitis (IPV) For OPV see page 2	4 doses The final dose of polio must be received at/after 4 years		

## ARIZONA GUIDE TO IMMUNIZATIONS REQUIRED FOR ENTRY - Minimum Interval/Catch-up Guidance; Grades K2 (School year 2023/2024)

Vaccine	Dose #	Minimum Age	Minimum Interval Between Doses	Notes
HepB Hepatitis B	dose 1	Birth	At least 4 weeks between dose 1 & 2	Some children may receive a birth dose and then a combination vaccine resulting in a total of 4 (or more) doses. As long as the interval between doses is met, 4+ doses meet requirements 2 doses, at least 4 months apart, meet the requirement if the child received the adolescent series using the Merck Recombivax HB Adult Formulation when the child was 11 years of age.
	dose 2	4 weeks	At least 8 weeks between dose 2 & 3 (or final)	
	dose 3	24 weeks	At least 16 weeks between dose 1 & (or final) AND at/after 24 weeks of age	
Polio IPV or OPV	dose 1 dose 2	6 weeks	At least 4 weeks between dose 1 & 2	Retrospectively: 1) A final dose given on or after August 7, 2009, must be given at or after 4 years of age and minimum interval of 6 months from the previous dose. 2) Students who received 4 doses (with at least 4 weeks minimum intervals between doses and/or before the age of 4 years) PRIOR to August 7, 2009 have met the requirement. OPV given prior to April 1, 2016 will be presumed to be trivalent and therefore acceptable, regardless of age, or country, of administration. Any OPV doses administered on or after April 1, 2016 are presumed to be bivalent and therefore unacceptable. Poliomyelitis vaccine is not recommended in the U.S. for individuals 18 years of age or older; however, a complete series is still required for school attendance.

Vaccine	Dose #	Minimum Age	Minimum Interval Between Doses	Notes
DTaP, Tdap, Td Tetanus, Diphtheria, and Pertussis	dose 1	6 weeks	At least 4 weeks between dose 1 & 2	<p>DTaP is licensed for children through age 6. If catch doses are needed at age 7 or older, Tdap or Td should be used to start/complete the series.</p> <p>A Tdap given at age 9 years of age does not count for the 11-12-year-old Tdap requirement; a Tdap should be given once 5 years has passed since the last dose of tetanus/diphtheria containing vaccines was given.</p> <p>Retrospectively, if a child received a Tdap at age 10 as part of a catch-up series, or inadvertently earlier than the recommended age of 11-12, the dose may be counted as the adolescent dose and is acceptable to meet school requirements.</p> <p>Once a valid adolescent Tdap dose has been received, a tetanus booster is recommended when 10 years has passed since last dose of tetanus-containing vaccine</p> <p>Refer to DTaP, Tdap, Td (Diphtheria, Tetanus, Pertussis) Grades K-12 Flowcharts</p>
	dose 2	10 weeks	At least 4 weeks between dose 2 & 3	
	dose 3	14 weeks	At least 6 months between dose 3 & 4	
	dose 4	12 months	At least 6 months between dose 4 & 5	
	dose 5	4 years		